arXiv:1410.7075v1 [math.CA] 5 Oct 2014

A NOTE ON THE VILENKIN-FOURIER COEFFICIENTS

GEORGE TEPHNADZE

Abstract. The main aim of this paper is to find the estimation for Vilenkin-Fourier coefficients.

2000 Mathematics Subject Classification. 42C10.

Key words and phrases: Vilenkin system, Fourier coefficients, martingale Hardy space.

Let P_+ denote the set of the positive integers, $P := P_+ \cup \{0\}$.

Let $m := (m_0, m_{1,...})$ denote a sequence of the positive integers not less than 2.

Denote by

$$Z_{m_k} := \{0, 1, ...m_k - 1\}$$

the additive group of integers modulo m_k.

Define the group G_m as the complete direct product of the group Z_{m_j} with the product of the discrete topologies of Z_{m_j} 's.

The direct product μ of the measures

$$\mu_k(\{j\}) := 1/m_k, \qquad (j \in Z_{m_k})$$

is the Haar measure on G_m with $\mu(G_m) = 1$.

If $\sup_n m_n < \infty$, then we call G_m a bounded Vilenkin group. If the generating sequence m is not bounded then G_m is said to be an unbounded Vilenkin group. **In this paper we discuss bounded Vilenkin groups only.**

The elements of G_m represented by sequences

$$x := (x_0, x_{1,...}, x_{j,...}), \qquad (\ x_k \in Z_{m_k}).$$

It is easy to give a base for the neighborhood of G_m

$$I_0(x) := G_m,$$

$$I_n(x) := \{y \in G_m \mid y_0 = x_0, ...y_{n-1} = x_{n-1}\}, \quad (x \in G_m,\ n \in P).$$

Denote $I_n := I_n(0)$ for $n \in P$ and $\overline{I_n} := G_m \setminus I_n$.

It is evident

$$\overline{I_N} = \bigcup_{s=0}^{N-1} I_s \setminus I_{s+1}. \tag{1}$$

If we define the so-called generalized number system based on m in the following way :

$$M_0 := 1, \qquad M_{k+1} := m_k M_k \, , \qquad (k \in P),$$

then every $n \in P$ can be uniquely expressed as $n = \sum_{k=0}^{\infty} n_j M_j$, where $n_j \in Z_{m_j}$ $(j \in P)$ and only a finite number of n_j's differ from zero. Let $|n| := \max \{j \in P;\, n_j \neq 0\}$.

Denote by $L_1(G_m)$ the usual (one dimensional) Lebesque space.

Next, we introduce on G_m an ortonormal system which is called the Vilenkin system.

At first define the complex valued function $r_k(x) : G_m \to C$, the generalized Rademacher functions as

$$r_k(x) := \exp(2\pi i x_k / m_k), \qquad \left(i^2 = -1,\ x \in G_m,\ k \in P\right).$$

Now define the Vilenkin system $\psi := (\psi_n : n \in P)$ on G_m as:

$$\psi_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}(x), \qquad (n \in P).$$

Specifically, we call this system the Walsh-Paley one if m=2.

The Vilenkin system is ortonormal and complete in $L_2(G_m)$ [1, 7].

Now we introduce analogues of the usual definitions in Fourier-analysis. If $f \in L_1(G_m)$ we can establish the the Fourier coefficients, the partial sums, the Dirichlet kernels with respect to the Vilenkin system in the usual manner:

$$\begin{aligned} \widehat{f}(k) &:= \int_{G_m} f \overline{\psi}_k d\mu, \ (\ k \in P\ ), \\ S_n f &:= \sum_{k=0}^{n-1} \widehat{f}(k)\, \psi_k, (\ n \in P_+,\ S_0 f := 0), \\ D_n &:= \sum_{k=0}^{n-1} \psi_n, (\ n \in P_+\ ). \end{aligned}$$

Recall that (see [1])

$$D_{M_n}(x) = \begin{cases} M_n\,, \text{ if } x \in I_n, \\ 0\,, \text{ if } x \notin I_n. \end{cases} \tag{2}$$

and

$$D_n(x) = \psi_n(x) \left(\sum_{j=0}^{\infty} D_{M_j}(x) \sum_{u=m_j-n_j}^{m_j-1} r_j^u(x) \right). \tag{3}$$

The norm (or quasinorm) of the space $L_p(G_m)$ is defined by

$$\|f\|_p := \left(\int_{G_m} |f|^p \, d\mu \right)^{1/p} \qquad (0 < p < \infty).$$

The $\sigma-$algebra generated by the intervals $\{I_n(x) : x \in G_m\}$ will be denoted by F_n $(n \in P)$. The conditional expectation operators relative to $F_n\,(n \in P)$ are denoted by E_n. Then

$$E_n f(x) = S_{M_n} f(x) = \sum_{k=0}^{M_n-1} \widehat{f}(k)\, w_k = |I_n(x)|^{-1} \int_{I_n(x)} f(x) d\mu(x),$$

where $|I_n(x)| = M_n^{-1}$ denotes the length of $I_n(x)$.

A sequence $F = \left(f^{(n)},\ n \in P\right)$ of functions $f^{(n)} \in L_1(G)$ is said to be a dyadic martingale if (for details see e.g. [8])

(i) $f^{(n)}$ is F_n measurable for all $n \in P$,

(ii) $E_n f^{(m)} = f^{(n)}$ for all $n \leq m$.

The maximal function of a martingale f is denoted by

$$f^* = \sup_{n \in P} \left| f^{(n)} \right|.$$

In case $f \in L_1$, the maximal functions are also be given by

$$f^*(x) = \sup_{n \in P} \frac{1}{|I_n(x)|} \left| \int_{I_n(x)} f(u)\, \mu(u) \right|.$$

For $0 < p < \infty$ the Hardy martingale spaces $H_p\ (G_m)$ consist of all martingales for which

$$\|f\|_{H_p} := \|f^*\|_{L_p} < \infty.$$

If $f \in L_1$, then it is easy to show that the sequence $(S_{M_n} f : n \in P)$ is a martingale. If $f = \left(f^{(n)}, n \in P\right)$ is martingale then the Vilenkin-Fourier coefficients must be defined in a slightly different manner:

$$\widehat{f}(i) := \lim_{k \to \infty} \int_{G_m} f^{(k)}(x)\, \overline{\Psi}_i(x)\, d\mu(x). \tag{4}$$

The Vilenkin-Fourier coefficients of $f \in L_1(G_m)$ are the same as those of the martingale $(S_{M_n} f : n \in P)$ obtained from f .

A bounded measurable function a is p-atom, if there exist a interval I, such that

$$\begin{cases} a) & \int_I a d\mu = 0, \\ b) & \|a\|_\infty \leq \mu(I)^{-1/p}, \\ c) & \operatorname{supp}(a) \subset I. \end{cases} \tag{5}$$

The Hardy martingale spaces $H_p(G)$ for $0 < p \leq 1$ have an atomic atomic characterization (see [8]):

Theorem W 1. A martingale $f = \left(f^{(n)}, n \in P\right)$ is in $H_p\,(0 < p \leq 1)$ if and only if there exist a sequence $(a_k, k \in P)$ of p-atoms and a sequence $(\mu_k, k \in P)$ of a real numbers such that for every $n \in P$

$$f=\sum_{k=0}^{\infty}\mu_k S_{M_n}a_k=f^{(n)}, \tag{6}$$

$$\sum_{k=0}^{\infty}|\mu_k|^p<\infty.$$

Moreover, $\|f\|_{H_p}\backsim\inf\left(\sum_{k=0}^{\infty}|\mu_k|^p\right)^{1/p}$, where the infimum is taken over all decomposition of f of the form (6).

It follows that if any operator is uniformly bounded on the p-atom, then it is a bounded operator from the Hardy space H_p to the space L_p. Moreover, the following theorem is true (see [9]):

Theorem W 2. Suppose that an operator T is sublinear and for some $0<p\leq 1$

$$\int\limits_{\overline{I}}|Ta|^p\,d\mu\leq c_p<\infty,$$

for every p-atom a ,where I denote the support of the atom. If T is bounded from L_∞ to L_∞ ,then

$$\|Tf\|_{L_p(G_m)}\leq c_p\|f\|_{H_p(G_m)}.$$

The classical inequality of Hardy type is well known in the trigonometric as well as in the Vilenkin-Fourier analysis. Namely,

$$\sum_{k=1}^{\infty}\frac{\left|\widehat{f}(k)\right|}{k}\leq c\|f\|_{H_1},$$

where the function f belongs to the Hardy space H_1 and c is an absolute constant. This was proved in the trigonometric case by Hardy and Littlewood [5] (see also Coifman and Weiss [4]) and for Walsh system in [6].

Weisz [8, 10] generalized this result for Vilenkin system and proved:

$$\sum_{k=1}^{\infty}\frac{|f(k)|^p}{k^{2-p}}\leq c\|f\|_{H_p}^p, \tag{7}$$

for all $f\in H_p$ $(0<p\leq 2)$.

It is also well-known (see [1]) that

$$\widehat{f}(n)\to 0,\ \text{when } n\to\infty,$$

for all $f\in L_1(G_m)$, where $\widehat{f}(n)$ denotes n-th Fourier coefficients of the function f.

The main aim of this paper is to prove that the following theorem is true:

Theorem 1. *a) Let $0<p<1$ and $f\in H_p(G_m)$. Then there exists an absolute constant c_p, defend only p, such that*

$$\left|\widehat{f}(n)\right| \leq c_p n^{1/p-1} \|f\|_{H_p}.$$

b) Let $0 < p < 1$ and $\Phi(n)$ is any nondecreasing, nonnegative function, satisfying condition

$$\overline{\lim_{n\to\infty}} \frac{n^{1/p-1}}{\Phi(n)} = \infty,$$

then there exists a martingale $f_0 \in H_p(G_m)$, such that

$$\overline{\lim_{n\to\infty}} \frac{\left|\widehat{f_0}(n)\right|}{\Phi(n)} = \infty.$$

Proof of Theorem 1. At first we prove that the maximal operator

$$\widetilde{S}_p^*(f) := \sup_{n\in P} \frac{|S_n f|}{(n+1)^{1/p-1}}$$

is bounded from the Hardy space H_p to the space L_p for $0 < p < 1$. Since the maximal operator $\widetilde{S}_p^*$ is bounded from L_∞ to L_∞ by Theorem W 2 we obtain that the proof of theorem will be complete, if we show that

$$\int_{\overline{I}_N} \left(\sup_{n\in P} \frac{|S_n a|}{(n+1)^{1/p-1}}\right)^p d\mu \leq c < \infty, \qquad \text{when } 0 < p < 1,$$

for every p-atom a $(0 < p \leq 1)$, where I denote the support of the atom. Let a be an arbitrary p-atom with support I and $\mu(I) = M_N^{-1}$. We may assume that $I = I_N$. It is easy to see that $S_n(a) = 0$ when $n \leq M_N$. Therefore we can suppose that $n > M_N$.

Let $x \in I_l \backslash I_{l+1}$. Combining (2) and (3) we have

$$|D_n(x)| \leq \sum_{j=0}^{l} n_j D_{M_j}(x) = \sum_{j=0}^{l} n_j M_j \leq c M_l. \tag{8}$$

Since $t \in I_N$ and $x \in I_s \backslash I_{s+1}$, $s = 0, \ldots N-1$, we obtain that $x - t \in I_s \backslash I_{s+1}$. Using (8) we get

$$|D_n(x-t)| \leq c M_s,$$

and

$$\int_{I_N} |D_n(x-t)|\, d\mu(t) \leq c\ \frac{M_s}{M_N}. \tag{9}$$

Hence

$$
\begin{aligned}
(10) \qquad & \frac{|S_n(a)|}{(n+1)^{1/p-1}} \\
\leq\ & \int_{I_N} |a(t)| \left| \frac{D_n(x-t)}{(n+1)^{1/p-1}} \right| d\mu(t) \\
\leq\ & \frac{\|a\|_\infty}{(n+1)^{1/p-1}} \int_{I_N} |D_n(x-t)|\, d\mu(t) \\
\leq\ & \frac{M_N^{1/p}}{(n+1)^{1/p-1}} \int_{I_N} |D_n(x-t)|\, d\mu(t) \\
\leq\ & \frac{cM_N^{1/p}}{M_N^{1/p-1}} \frac{M_s}{M_N} = cM_s.
\end{aligned}
$$

Combining (1) and (10) we have

$$
\begin{aligned}
& \int_{\overline{I_N}} \left| \widetilde{S}_p^* a(x) \right|^p d\mu(x) \\
=\ & \sum_{s=0}^{N-1} \int_{I_s \setminus I_{s+1}} \left| \widetilde{S}_p^* a(x) \right|^p d\mu(x) \\
\leq\ & c \sum_{s=0}^{N-1} \frac{M_s^p}{M_s} < c < \infty.
\end{aligned}
$$

Now we are ready to prove the main result. Let $0 < p < 1$.Then

$$
\left| \widehat{f}(n) \right| = |S_{n+1}f - S_n f| \leq 2 \sup_{n \in P} |S_n f|
$$

and

$$
(11) \qquad \frac{\left| \widehat{f}(n) \right|}{(n+1)^{1/p-1}} \leq 2 \sup_{n \in P} \frac{|S_n f|}{(n+1)^{1/p-1}},
$$

Consequently,

$$
\frac{\left| \widehat{f}(n) \right|}{(n+1)^{1/p-1}} \leq 2 \left\| \sup_{n \in P} \frac{|S_n f|}{(n+1)^{1/p-1}} \right\|_p \leq c_p \|f\|_{H_p}.
$$

It follows that

$$
\left| \widehat{f}(n) \right| \leq c_p n^{1/p-1} \|f\|_{H_p}.
$$

b) In the proof of second part of theorem we follow the method of Blahota, Gát and Goginava (see [2, 3]).

Let $0 < p < 1$ and $\Phi(n)$ is any nondecreasing, nonnegative function, satisfying condition

$$\overline{\lim_{n\to\infty}} \frac{n^{1/p-1}}{\Phi(n)} = \infty, \tag{12}$$

then for every $0 < p < 1$, there exists an increasing sequence $\{\alpha_k : k \in P\}$ of the positive integers such that:

$$\sum_{k=0}^{\infty} \left(\frac{\Phi(M_{\alpha_k})}{M_{\alpha_k}^{1/p-1}} \right)^{p/2} < \infty \tag{13}$$

Let

$$f^{(A)}(x) = \sum_{\{k;\ \alpha_k < A\}} \lambda_k a_k,$$

where

$$\lambda_k = \left(\frac{\Phi(M_{\alpha_k})}{M_{\alpha_k}^{1/p-1}} \right)^{1/2}$$

and

$$a_k(x) := \frac{M_{\alpha_k}^{1/p-1}}{M} \left(D_{M_{\alpha_k+1}}(x) - D_{M_{\alpha_k}}(x) \right).$$

It is easy to show that the martingale $f = \left(f^{(1)}, f^{(2)} ... f^{(A)} ...\right) \in H_p$.

Indeed, since

$$S_{M_A} a_k(x) = \begin{cases} a_k(x), & \alpha_k < A, \\ 0, & \alpha_k \geq A, \end{cases} \tag{14}$$

$$\operatorname{supp}(a_k) = I_{\alpha_k},$$

$$\int_{I_{\alpha_k}} a_k d\mu = 0$$

and

$$\|a_k\|_\infty \leq \frac{M_{\alpha_k}^{1/p-1}}{M} M_{\alpha_k+1} \leq (M_{\alpha_k})^{1/p} = (\operatorname{supp}\ a_k)^{-1/p}$$

if we apply Theorem W 1 and (13) we conclude that $f \in H_p$.

It is easy to show that

$$\widehat{f}(j) = \begin{cases} \frac{1}{M} M_{\alpha_k}^{(1/p-1)/2} \Phi^{1/2}(M_{\alpha_k}), & \text{if } j \in \{M_{\alpha_k}, ..., M_{\alpha_k+1} - 1\},\ k = 0, 1, 2... \\ 0, & \text{if } j \notin \bigcup_{k=1}^{\infty} \{M_{\alpha_k}, ..., M_{\alpha_k+1} - 1\}. \end{cases} \tag{15}$$

It follows that

$$\begin{aligned}\overline{\lim_{n\to\infty}}\frac{\widehat{f}(n)}{\Phi(n)} &\geq \lim_{k\to\infty}\frac{\widehat{f}(M_{\alpha_k})}{\Phi(M_{\alpha_k})}\\ &\geq \lim_{k\to\infty}\frac{1}{M}\frac{M_{\alpha_k}^{(1/p-1)/2}\Phi^{1/2}(M_{\alpha_k})}{\Phi(M_{\alpha_k})}\\ &\geq \lim_{k\to\infty}\left(\frac{M_{\alpha_k}^{(1/p-1)}}{\Phi(M_{\alpha_k})}\right)^{1/2}=\infty.\end{aligned}$$

Theorem 1 is proved.

References

[1] *G. N. Agaev, N. Ya. Vilenkin, G. M. Dzhafarly and A. I. Rubinshtein,* Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehim, 1981 (in Russian).

[2] *I. Blahota, G. Gát* and *U. Goginava,* Maximal operators of Fejer means of double Vilenkin-Fourier series, Colloq. Math. 107 (2007), no. 2, 287–296.

[3] *I. Blahota, G. Gát* and *U. Goginava,* Maximal operators of Fejér means of Vilenkin-Fourier series. JIPAM. J. Inequal. Pure Appl. Math. 7 (2006), 1- 7 .

[4] *R. Coifman, G. Weiss,* Extensions of the Hardy spaces and their use in analysis. Bull. Amer. Math. Soc. 83, (1977), 569-645.

[5] *G.H. Hardy, J.E. Littlewood,* Some new properties of Fourier constants. J. London Math. Soc. 6, (1931), 3-9.

[6] *F.Schipp, W.R. Wade, P. Simon. J. Pál,* Walsh series, An Introduction to Duadic Harmonic Analysis, Akademiai Kiadó, (Budapest-AdamHilger (Bristol-New-York)), 1990.

[7] *N. Ya. Vilenkin,* A class of complate ortonormal systems, Izv. Akad. Nauk. U.S.S.R., Ser. Mat., 11 (1947), 363-400.

[8] *F. Weisz,* Martingale Hardy spaces and their applications in Fourier Analysis, Springer, Berlin-Heideiberg-New York, 1994.

[9] *F. Weisz,* Summability of multi-dimensional Fourier series and Hardy space, Kluwer Academic, Dordrecht, 2002.

[10] *F. Weisz*, Hardy-Littlewood inequalities for Ciesielski-Fourier series, Analysis Mathematica, 31 (2005), 217-233.

G. Tephnadze, Department of Mathematics, Faculty of Exact and Natural Sciences, Tbilisi State University, Chavchavadze str. 1, Tbilisi 0128, Georgia

E-mail address: giorgitephnadze@gmail.com